





Opioid and Methamphetamine Use Disorders: Diagnosis and Treatment

Objectives



1. Review the diagnostic criteria for substance use disorders
2. Overview of the neurobiology of addiction
3. Review office-based medications to treat opioid use disorder
4. Discuss interventions to treat methamphetamine use disorder

Objectives



- 1. Review the diagnostic criteria for substance use disorders**

DSM 5

11 criteria

Diagnostic and Statistical Manual of
Mental Disorders


4 C's

Craving

Compulsion


Consequences

Loss of Control




DSM 5: Substance Use Disorder

- Taking in larger amounts or for longer than intended
- Unsuccessful efforts to cut down
- Spending a lot of time obtaining the substance
- Craving or a strong desire to use the substance



DSM 5: Substance Use Disorder

- Continued use despite recurring social or interpersonal problems due to use
- Important activities given up or reduced
- Recurrent use in physically hazardous situations
- Persistent / Recurrent physical or psychological difficulties from use
- Recurrent use resulting in a failure to fulfill major role obligations




DSM 5: Substance Use Disorder



Tolerance*



Withdrawal*



Substance Use Disorder

2–3

mild disorder

4–5

moderate disorder

6+

severe disorder

Substance Use Disorder



Substance Use Disorder

Diagnosis



Addict

Label/Accusation

The words we use to describe our patients affects the care they receive

Substance Use Disorder


Recovery Dialects
The words we use matter.

Positive		Negative
Person who uses substances		Substance Abuser
Recurrence of Use		Relapse
Pharmacotherapy		Medication-Assisted Treatment
Accidental Drug Poisoning		Overdose
Person with a Substance Use Disorder		Addict
		Alcoholic
		Opioid Addict

While some negative language is okay to use in mutual aid meetings, its use should be avoided in public, when advocating and in journalism.


SOURCE: Ashford, R. D., Brown, A. M., & Curtis, B. (2018). Substance use, recovery, and linguistics: The impact of word choice on explicit and implicit bias. *Drug and Alcohol Dependence*, 189, 131–138.

Objectives



2. Overview of the neurobiology of addiction


Mesolimbic Dopamine System




1950s : electrodes



1970s: Dopamine




Desire and Drives




The use of dopamine neurons to shape responses to rewards is seen in simple organisms like worms and flies.

It evolved millions of years ago.


Dopaminergic impulses tell organisms to move toward reward (warmth, food, moisture)



- In humans, those dopaminergic impulses travel through the NAC
- Mediates responses to food, sex, social interactions
- DA projections from VTA to NA release DA and tell the NA to go for it!
- Connects with memory and emotional centers so it can be repeated in the future




- Addiction taps into this normal brain process
- All addictive drugs activate this pathway
- Drug experience is deeply linked to memory and emotion




- People, places, things associated with drug use can trigger cravings
- Even when images associated with drug use are shown too rapidly to be “seen” they still trigger cravings

Dysregulation



- Dysregulation: impaired ability of the front of the brain to regulate what is going on in the older regions of the brain.
- Prefrontal cortex helps determine the risks and benefits of behaviors and make rational choices.



- Prefrontal cortex is newer and more complicated. It needs a little time to weigh in.
- Repeated activation of the VTA to NAC track slowly strengthens those connections. Habits get hard wired, fast and automatic



Little pleasures like family, friends, jobs well done, tasks accomplished, provide just enough dopamine to activate the D2 receptors and strengthen the impulses that slow things down.


Medications to decrease craving, attenuate withdrawal symptoms, and decrease deaths

Behavioral interventions that entrain different habits

Conclusion

- Addiction taps into normal brain processes
- It is entrained through habit
- It can be effectively treated

Objectives




3. Review office-based medications to treat opioid use disorder


What are they?

Methadone
Buprenorphine
XR- Naltrexone

Pharmacotherapy for Opioid Use Disorder



Pharmacotherapy for Opioid Use Disorder




Why do they matter now more than ever?



Fentanyl




Fentanyl



All Overdose Deaths 1999-2020

Overdose Deaths by Drug 1999-20



(source)

** 2020 numbers are reported provisional deaths per CDC, subject to change

Fentanyl



High affinity and high efficacy at mu receptor

Single use has a short half-life (fast on, fast off)

Repeated use may lead to accumulation in adipose tissue, decreased renal clearance, more mu opioid receptor desensitization

Buprenorphine



Why is it so great?


It decreases opioid
cravings, withdrawal, and
use.



Patients taking buprenorphine are significantly more likely to engage and remain in treatment compared to those tapered off the medication.

Fiellen 2014; D'Onofrio 2017


Buprenorphine: Maintenance vs. Taper



Why is it so great?

Most importantly,
people don't die

Mortality Risk during and after buprenorphine treatment




Mortality Risk during and after opioid substitution treatment: systematic review and meta-analysis of cohort studies. Sordo, et al. BMJ 2017.

Important to know:



- Buprenorphine is a high affinity binder at the mu opioid receptors. That means it sits tightly on the receptor.
- It will kick off anything else that's bound there
- But it is a partial agonist at the receptor. That means it doesn't activate the receptor completely.
- If it kicks a full agonist off the receptor, the difference between full agonism and partial agonism is big enough \square precipitated withdrawal


Available in two primary forms:



1. Buprenorphine monoprodut (Subutex)
2. Buprenorphine/Naloxone (Suboxone)

Buprenorphine/naloxone may reduce misuse

- Buprenorphine is taken sublingually
- Naloxone is absorbed in minute amounts sublingually.
- It is essentially inactive (in most people) unless injected
- Decreased risk of misuse (controversial)



Newer kids in town: buprenorphine XR (Sublocade and Brixadi)



Sublocade:

Approved November 2017

Single injection lasts one month

Must be refrigerated

Injection hurts

Brixadi:

Approved May 2023

Single injection lasts 1 week or 1 month, various doses

Does not require refrigeration

Injection not as painful



How to administer and prescribe

“Traditional” inductions



- Instruct the patient to abstain from any opioid use for a minimum of:
 - 12-16 hours for short-acting opioids
 - 24 hours for sustained-release opioid medications
 - 36 hours for methadone or fentanyl
- Observe and document mild to moderate withdrawal

“Traditional” inductions



Wait until patient is in mild to moderate withdrawal (which means receptors are empty)

Begin buprenorphine and titrate up, as needed, over 3-4 days



How do you know if a patient is in sufficient enough withdrawal to begin buprenorphine?

Clinical Opiate Withdrawal Scale (COWS)


Clinical Opiate Withdrawal Scale

For each item, circle the number that best describes the patient's signs or symptom. Rate on just the apparent relationship to opiate withdrawal. For example, if heart rate is increased because the patient was jogging just prior to assessment, the increase pulse rate would not add to the score.

Patient's Name: _____		Date and Time ____/____/____ : ____:____
Reason for this assessment: _____		
Resting Pulse Rate: _____ beats/minute <i>Measured after patient is sitting or lying for one minute</i> 0 pulse rate 80 or below 1 pulse rate 81-100 2 pulse rate 101-120 4 pulse rate greater than 120	GI Upset: over last ½ hour 0 no GI symptoms 1 stomach cramps 2 nausea or loose stool 3 vomiting or diarrhea 5 Multiple episodes of diarrhea or vomiting	
Sweating: over past ½ hour not accounted for by room temperature or patient activity. 0 no report of chills or flushing 1 subjective report of chills or flushing 2 flushed or observable moistness on face 3 beads of sweat on brow or face 4 sweat streaming off face	Tremor observation of outstretched hands 0 No tremor 1 tremor can be felt, but not observed 2 slight tremor observable 4 gross tremor or muscle twitching	
Restlessness Observation during assessment 0 able to sit still 1 reports difficulty sitting still, but is able to do so 3 frequent shifting or extraneous movements of legs/arms 5 Unable to sit still for more than a few seconds	Yawning Observation during assessment 0 no yawning 1 yawning once or twice during assessment 2 yawning three or more times during assessment 4 yawning several times/minute	
Pupil size 0 pupils pinned or normal size for room light 1 pupils possibly larger than normal for room light 2 pupils moderately dilated 5 pupils so dilated that only the rim of the iris is visible	Anxiety or Irritability 0 none 1 patient reports increasing irritability or anxiousness 2 patient obviously irritable anxious 4 patient so irritable or anxious that participation in the assessment is difficult	
Bone or Joint aches If patient was having pain previously, only the additional component attributed to opiates withdrawal is scored 0 not present 1 mild diffuse discomfort 2 patient reports severe diffuse aching of joints/ muscles 4 patient is rubbing joints or muscles and is unable to sit still because of discomfort	Gooseflesh skin 0 skin is smooth 3 piloerection of skin can be felt or hairs standing up on arms 5 prominent piloerection	
Runny nose or tearing Not accounted for by cold symptoms or allergies 0 not present 1 nasal stuffiness or unusually moist eyes 2 nose running or tearing 4 nose constantly running or tears streaming down cheeks	<p style="text-align: right;">Total Score _____</p> <p style="text-align: center;">The total score is the sum of all 11 items</p> Initials of person completing Assessment: _____	

Score: 5-12 = mild; 13-24 = moderate; 25-36 = moderately severe; more than 36 = severe withdrawal

Clinical Opiate Withdrawal Scale (COWS)

- 
- Resting pulse rate
 - Sweating/chills
 - Restlessness
 - Pupil size
 - Bone or joint aches
 - Runny nose
 - GI upset
 - Tremor
 - Yawning
 - Anxiety or irritability
 - Goose bumps
- **Guides timing of first dose of buprenorphine**

Traditional induction



Begin buprenorphine with COWS is 10-12

Prepare for Discomfort

- Acetaminophen and ibuprofen
- Clonidine
- Hydroxyzine
- Trazodone
- Tizanidine or Methocarbamol
- Ondansetron
- Bismuth or Loperamide

Srivastara, 2020; Kosten, 2019;
Kuzmaul 2020; Kheirabadi 2008
;Salehi 2011; Sanders 2013

Begin the medication

Give initial dose of buprenorphine (originally 4 mg) and titrate up over a period of hours.



Precipitated Withdrawal



If opioid withdrawal appears shortly after the first dose buprenorphine may have precipitated a withdrawal syndrome

Precipitated Withdrawal



Greatest severity of buprenorphine-related precipitated withdrawal in the first few hours (1-4) after a dose

Challenges with Traditional Induction

- Patient must experience withdrawal, which is difficult
- With fentanyl, sometimes need to wait even longer than 3 days because fentanyl sticks around in the fat
- Always possible that patient will experience precipitated withdrawal



Another option...



ninja clipart PNG Designed By 588ku from
https://pngtree.com/freepng/sneak-attack-sneak-attack-man-in-black-black-man-ninja_3931511.html?sol=downref&id=bef

Low dose buprenorphine induction


- Many different protocols
 - Initial protocol “Bernese Method”
 - Usually start at 0.5 mg
 - Often 7-10 days
 - No universally accepted regimen
 - Can continue full agonists throughout the entire induction

Day	Dose
1	0.5 mg daily
2	0.5 mg bid
3	1 mg bid
4	2 mg bid
5	4 mg bid
6	4 mg tid
7	8 mg tid

Adapted from Yale protocol

⁹Opioid Use Disorder Practice Update (2022)
British Columbia Centre on Substance Use

Rapid low dose inductions



Day	Full Opioid Agonist	Buprenorphine Dosing Instructions	Total Daily Dose of Buprenorphine
1	Continue	0.5 mg SL once	0.5 mg
2	Continue	0.5 mg SL bid	1 mg
3	Continue	1 mg SL bid	2 mg
4	Continue	2 mg SL bid	4 mg
5	STOP (if able to tolerate increase)	4 mg SL once. If tolerated take additional 4 mg in 10 mins. Continue to titrate prn for ongoing cravings or withdrawal symptoms for TDD of 16-24 mg	16-24 mg

Tips and Tricks

- Specifically outline what adjunct meds you are giving and for what
- Instruct patients to take AM buprenorphine before their full agonist



Tips and Tricks

- Close follow up
- Modify or slow protocol as needed (i.e. repeat days)
- Give naloxone to every patient



Maintenance

- Continue patient at the dose at which they have no withdrawal symptoms and minimal to no cravings
- The maximum effective dose has long been considered 24mg
- However, with fentanyl, many patients continue to have cravings and withdrawal symptoms at typical doses (16-24 mg)
- It is becoming more common to **up titrate** to 28-32 mg which seems to be helpful for some patients



Regulations and Regulatory Changes




Buprenorphine approved by the FDA in 2002. Prescribers were required to undergo an 8-hour training, register with the DEA, obtain an “X-waiver” and could only prescribe to 30 patients at a time


2016 NPs and PAs were allowed to prescribe, but with a longer training requirement. Still required to obtain X waiver and register with the DEA and limit patients

Training requirement removed in 2021, though prescribers still needed to obtain the waiver and register with the DEA

Jan 2023 all buprenorphine specific DEA requirements were removed

Naltrexone for Extended Release Injectable Suspension





Difficult to start

Requires abstinence from opioids 4 – 7 days

About 25% of patients will not complete induction



Overdose data


- Original findings
 - more overdoses in the XR-NTX arm, but not statistically significant
- Re-analysis
 - Researchers had missed cases of overdose
 - 28 overdoses in XR-NTX arm
 - 2.4 x greater hazard of overdose compared to bup/nal

Summary

- Opioid use disorder can be treated in an outpatient setting
- Buprenorphine saves lives
- Please prescribe



Objectives



4. Discuss interventions to treat methamphetamine use disorder

Crystal Methamphetamine



- Form of d-methamphetamine
- Closely related to amphetamine
- Longer lasting and more toxic to the CNS



Methamphetamine Crystals
Photo by Barbadiison, © 2011 Erowid.org



Crystal Methamphetamine





Crystal Methamphetamine



Crystal Methamphetamine



Crystal Methamphetamine



Crystal Methamphetamine

Materials:

1 2 Liter Bottle (with cap)
1 1 Liter Bottle (get 2 caps for it)
1 20 oz. Bottle (with cap)
1 Quart Jar
2 ft. 1/4in. diameter rubber/plastic hose (aquarium hose works good)
Coffee Filters
1 Funnel
1 Tubing Cutter (go to Home Depot)
2 Plyers
1 Roll of Ductape or Electrical Tape 1 Blender or Food Processor

200 60mg Pseudophedrine HCL pills (Actifed, Sudafed, Suphedrine, etc.)
1 1/2 cups Ammonium Nitrate fertilizer (33-0-0)
3 cans starting fluid
3 AA Energizer Lithuim Batteries
1 bottle Red Devil brand Lye
2 caps of water (use the top off the 2 liter)
1 box Iodized Salt
1 bottle Liquid Fire brand drain opener

Procedure:

1) Rinse and dry out all of your bottles. Be sure to get ALL of the moisture out. Don't go any further until they are completely dry.
2) Put your pills into the blender or food processor and grind them into powder. Mix them in with the 1 1/2 cups of Ammonium Nitrate fertilizer. Use the funnel to pour the mixture into the 2 liter bottle.



Crystal Methamphetamine



2005: CMEA (Combat Methamphetamine Epidemic Act)




Result?




In Oregon, from 2004 to 2011, methamphetamine lab incidents decreased from an average of 24 per month to less than one per month

And Yet...









Increase dopamine to +/- 200 times basal output





Effects of Drugs on Dopamine Release



Di Chiara and Imperato, PNAS, 1988

Medical Issues Related to Methamphetamine Use



Neurotoxicity, cognitive effects

- Excessive DA damaging cell structures
- Disruption of blood-barrier
- Use associated with poorer performance on motor and processing tasks, visual and verbal fluency
- More than 2/3 of those with MUD show cognitive impairment
- May limit ability to follow through with treatment, understand advice, and achieve treatment outcomes

Medical Issues Related to Methamphetamine Use



Cardiovascular and cerebrovascular

- Leading cause of death with MUD
- Strokes more common in young men (hemorrhagic)
- Also associated with pulmonary htn, cardiac arrhythmia, cardiomyopathy

Lappin et al., 2017



Two evidence-based behavioral interventions: contingency management and harm reduction



- Behavioral interventions = first line treatment for MUD
- Most behavioral interventions (CBT, MI, Matrix model, exercise, CM) demonstrated some efficacy in reducing methamphetamine cravings and use
- Contingency management most consistently showed reduced use, increased retention in treatment, better quality of life



Photo courtesy of John Mahan MD

Contingency Management: Theory



- Addiction is sustained through reinforced learning
- We cannot simply unlearn habits – we must learn new and competing habits
- CM entrains new behaviors that support the process of recovery
- Breaks recovery process down into a series of concrete, attainable goals
- > 100 RCTs affirm the effectiveness of CM in treating addiction

Roll JM et al. Am Jnl Psych 2006
Roll JM et al. Addict Behav 2013
Rawson, RA et al. Addiction 2016

Contingency Management: Practice



1. Identify a target behavior that can be objectively measured, attainable, and reinforced in real time.
2. Reward that behavior immediately when it occurs, using rewards that are valuable to participants (but not necessarily expensive).
3. Use an escalating schedule of reinforcement.



Photo courtesy of John Mahan MD

Example



Patient on long term IV antibiotics who is often not in her room when it is time for her antibiotics. She likes chocolate and Starbuck's Frappuccinos

Target behavior: be in the room 8:00 am, noon, and 5 pm

Reward: Hershey's kiss each time she is in the room when the nurse arrives with antibiotics

Escalating schedule: \$5 Starbuck's card after she has accumulated 10 Hershey's kisses

Harm Reduction



Harm reduction is a set of practical strategies and ideas aimed at reducing negative consequences associated with drug use. Harm Reduction is also a movement for social justice built on a belief in, and respect for, the rights of people who use drugs.

Harm Reduction is also



Part of the continuum of care

Relationship building

Treatment

Harm Reduction is not



What we do when nothing else works

Harm Reduction Practices: Methamphetamines

Safe injecting:



- Clean needles/rigs (including don't share filters, cookers)
- Don't use alone
- Use needles bevel up
- Use a filter whenever possible
- Test for fentanyl
- Clean water

Collins S et al. *Intl Jnl of Drug Policy* 2019

Thakrar K, Weinstein ZM, Walley AY. *Postgrad Med J.* 2016;92(1088):356–363.

Harm Reduction Practices: Methamphetamines



Hydration

Toothbrushes

Condoms

Naloxone

Patient Centered: Ask the patient/client: what harms most concern you?

Meds for MA/A Use Disorder



- No FDA-approved meds for MA/A use disorder (MUD)
- Lots of research looking into possible treatments
- Will review published findings from 2 recent trials
- Systematic reviews of medications for MUD


Mirtazapine



- FDA-approved antidepressant
- Main side effects weight gain and somnolence
- Mixed monoamine agonist-antagonist
- Cisgender men & transgender women sex w/ men
- Double blind RCT of 120 participants
- ↓ in methamphetamine + UDT despite low adherence

Naltrexone IM + Bupropion

- Large multi-center RCT, two-stage, sequential parallel comparison design.
- Number needed to treat – 9, low treatment improvement



Sufficient Evidence of No Benefit



- Dopamine agonists (levodopa, cabergoline, pramipexole)
- Antipsychotics – aripiprazole
- Antidepressants – SSRIs
- Anticonvulsants/muscle relaxants
- Varenicline

Briones M et al. Drug Alcohol Depend 2018
Ronsley C et al PLoS ONE 2020
Chan B et al Addiction 2019

Insufficient Evidence of Benefit



- Prescription psychostimulant agonist therapy – methylphenidate, modafinil, lisdexamphetamine, dextroamphetamine, mixed amphetamine salts
- Antidepressants – non-SSRI (mirtazapine, bupropion)
- N-acetylcysteine (NAC) – acts as a physiological reservoir of neuronal glutamate


Coffin P et al JAMA Psychiatry 2020

Ronsley C et al PLoS ONE 2020

Tardelli VS et al Psychopharmacology 2020

Chang C-T et al Clin Psychopharmacol Neurosci 2021

Summary



- Methamphetamine use and use disorders are escalating
- There are effective behavioral interventions
- Harm reduction is treatment
- Medications are being investigated

- Questions?

- Thoughts?