

Indian Country Infectious Disease Update June 2024

Jorge Mera, MD, FACP

And

Jonathan Iralu, MD, MACP, FIDSA




Look out for Ocular syphilis...

- Quite common in the current outbreak
- Can involve any part of the eye
 - ❖Keratitis of the cornea
 - ❖Iritis (anterior uveitis) often seen with secondary syphilis
 - Vitritis: patient sees a starry sky, small white balls seen on exam
 - Choreoretinitis:
 - ground glass retinitis caused by syphilitic vasculitis
 - yellow plaques on retina near macula in Acute Posterior Placoid Chorioretinopathy
 - Optic neuritis: monocular vision loss


Ocular Syphilis

- Symptoms:
 - **❖**Floaters
 - ❖Eye pain/photophobia
 - Vision loss
- ❖ Findings:
 - Diverse depending on site of infection
 - Argyll-Robertson pupils are a marker for neurosyphilis, not ocular syphilis
 - ❖Irregular pupil
 - ❖React to accommodation, not light.





Syphilitic chorioretinitis



Ocular Syphilis Management

- ❖ Send all syphilis patients with eye complaints or findings to eye clinic STAT
- ❖ Treatment is identical to neurosyphilis
- ❖ LP is not required unless the neuro exam (especially cranial nerve exam) is abnormal
- ❖ Follow RPR titer and eye exam after treatment

Doxy PEP is the prevention tool

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Postexposure Doxycycline to Prevent Bacterial Sexually Transmitted Infections

Anne F. Luetkemeyer, M.D., Deborah Donnell, Ph.D.,
Julia C. Dombrowski, M.D., M.P.H., Stephanie Cohen, M.D., M.P.H.,
Cole Grabow, M.P.H., Clare E. Brown, Ph.D., Cheryl Malinski, B.S.,
Rodney Perkins, R.N., M.P.H., Melody Nasser, B.A., Carolina Lopez, B.A.,
Eric Vittinghoff, Ph.D., Susan P. Buchbinder, M.D., Hyman Scott, M.D., M.P.H.,
Edwin D. Charlebois, Ph.D., M.P.H., Diane V. Havlir, M.D., Olusegun O. Soge, Ph.D.,
and Connie Celum, M.D., M.P.H., for the DoxyPEP Study Team*

Luetkemeyer et al NEJM, 4/6/2023

- Recruited 501 MSM and transgender women on PrEP or ART
- Offered Doxycycline 200 mg within 72 hours of condomless sex
- ❖ Relative risk for infection

		PrEP	ART
*	Gonorrhea	0.45	0.43
*	Chlamydia	0.12	0.26
*	Syphilis	0.13	0.23


❖ Adverse events: None serious

https://www.nejm.org/doi/pdf/10.1056/NEJMoa2211934?articleTools=true

Doxycycline Post Exposure Prophylaxis New CDC guideline!

- ❖ Patients with STI risk can be offered 200 mg doxycycline within 24-72 hours of sex
 - ❖ MSM
 - ❖ Bisexual
 - ❖ Transgender women
- Indicated for
 - ❖ Bacterial STI within the last year
 - Participating in sexual activities that are known to increase the likelihood of exposure to STIs (see CDC fine print).

https://www.cdc.gov/mmwr/volumes/73/rr/pdfs/rr7302a1-H.pdf





Expanded Testing Panel

January 25, 2024

Tests

Chlamydia – genital, oral, rectal

Gonorrhea – genital, oral, rectal

Trichomonas – vaginal

HIV (New)


Syphilis (New)

Hepatitis B (New)

Hepatitis C (New)



Provided Materials




Sample Quantities

- 450 μL minimum required
- 650 μL preferred


Procedure













Marketing







Ordering

- Attention IWTK Users: We have added a new feature that allows you to have two separate
 addresses: your physical (home) address and your mailing address. If these two addresses are
 the same, you may indicate that in your IWTK account. You may update either address on file at
 any time in your IWTK account.
- If you are not a resident of MD, OK, KS, AK, NM, UT or AZ and would like to be tested for STIs, please visit the CDC Testing Locator to find a testing site near you.

