



**YOUR GUIDE TO GIVING
MAVYRET TO YOUR CHILD**

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C):

- Genotypes (GT) 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis
- Or GT 1 infection and have been previously treated with a regimen that contained a hep C NS5A inhibitor or an NS3/4A protease inhibitor, but not both.

SAFETY CONSIDERATIONS

- **Hepatitis B virus (hep B) reactivation:** Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.
- Do not take MAVYRET if you have certain liver problems or if you are taking the medicines atazanavir or rifampin.

Please see additional Important Safety Information throughout the brochure, including **Boxed Warning**.

Please see accompanying full **Prescribing Information**, including the **Patient Information**.

YOU'VE MADE A BIG STEP TOWARD TREATING YOUR CHILD'S HEPATITIS C.

MAVYRET oral pellets should be given as instructed by your child's healthcare provider.

The next page includes step-by-step instructions on how to prepare MAVYRET. It may be helpful to keep these instructions near the medication during your child's treatment.

Have additional questions about taking MAVYRET?
Call 877-MAVYRET (877-628-9738)

SAFETY CONSIDERATIONS

- In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death. Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.
- There are some medicines that you should not take with MAVYRET. **Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work or may cause side effects. **Do not start a new medicine without telling your doctor.**

Please see additional Important Safety Information throughout the brochure, including **Boxed Warning**.

Please see accompanying full **Prescribing Information**, including the **Patient Information**.



READ STEPS 1 - 5 before you get started!

1
Find the number of packets



Mix the day's packets together with food.

About 1-2 teaspoons of an acceptable food like the ones below

Remember, if you need help, you can always call!
Call your healthcare provider, your pharmacist, or us: 877-MAVYRET (877-628-9738)

2
Choose the right food



IMPORTANT: If the pellets are chewed, or dissolve in the food, the medicine tastes bad and might not work.




YES
Choose soft foods that stick to the spoon





NO
Watery, heated, frozen, or solid foods

3
Gather what you'll need




4
Prepare the dose



MIX
thoroughly (don't crush, or it will taste bitter)

5
Give the medication



! QUESTIONS AND SAFETY INFORMATION

How long can the pellets stay in the mixture?
• After 5 minutes, a bitter taste will be noticeable to the child.
• After 15 minutes, the medication will be less effective.
Discard and call your healthcare provider.

What if my child missed a daily dose of the medication?
• Contact your healthcare provider. **Do not** give a double dose of MAVYRET.

What if my child is allergic to the recommended foods?
• Contact your healthcare provider. **Do not** mix with food that your child is allergic to.

What if my child has trouble swallowing or cannot finish the mixture?
• Contact your healthcare provider. **Do not** continue giving the mixture or give your child an additional dose of MAVYRET.

Can my child drink water after completing their dose?
• Yes. Drinking water after taking the medication will not change its effectiveness.

What if I need replacement packets?
• Contact your healthcare provider.

Please see additional Important Safety Information throughout the brochure, including Boxed Warning. Please see accompanying full Prescribing Information, including the Patient Information.

NEED HELP?



Call your healthcare provider, call your pharmacist, call us at **877-MAVYRET (877-628-9738)** or visit www.mavyret.com

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MAVYRET?

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

Do not take MAVYRET if you:

- Have certain liver problems
- Are taking the medicines atazanavir or rifampin

What should I tell my doctor before taking MAVYRET?

- If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.
- If you are pregnant or plan to become pregnant, or if you are breastfeeding or plan to breastfeed. It is not known if MAVYRET will harm your unborn baby or pass into your breast milk. Talk to your doctor about the best way to feed your baby if you take MAVYRET.
- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work or may cause side effects.
- **Do not start taking a new medicine without telling your doctor.** Your doctor can tell you if it is safe to take MAVYRET with other medicines.

What are the possible side effects of MAVYRET?

- **In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death.** Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.
- The most common side effects of MAVYRET are headache and tiredness.

These are not all the possible side effects of MAVYRET. Call your doctor for medical advice about side effects.

This is the most important information to know about MAVYRET. For more information, talk to your doctor or healthcare provider.

MAVYRET oral pellets are dispensed in unit-dose packets. Each packet contains 50 mg glecaprevir/20 mg pibrentasvir.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying full **Prescribing Information**, including the **Patient Information**.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Visit MAVYRET.com for more information about hep C treatment

abbvie

©2022 AbbVie Inc. North Chicago, IL 60064
US-MAVY-220119 April 2022

MAVYRET 
glecaprevir/pibrentasvir
100 mg/40 mg tablets
50 mg/20 mg oral pellets